


AGU Fall Meeting 2009

You may print by clicking on this  button. To return to the previous page, close this browser window or click the 'X' button in the top right corner of the page.

ID# GC31A-0695

Location: Poster Hall (Moscone South)

Time of Presentation: Dec 16 8:00 AM - 12:20 PM

Climate- and Anthropogenically-Induced Change in the Republic of Tyva

A. J. Soja^{1, 6}; *N. Tchebakova*²; *E. Parfenova*²; *V. Kanza*⁴; *D. J. Westberg*⁵; *L. Buryiak*³; *O. Kalenskaya*³; *E. A. Kukavskaya*²; *G. A. Ivanova*²; *P. W. Stackhouse*⁶; *A. Sukhinin*²

1. Climate Science and Radiation and Aerosols Branches, National Institute Aerospace, Hampton, VA, United States.

2. Sukachev Institute of Forestry, Krasnojarsk, Krasnojarsk, Russian Federation.

3. State Siberian Technological University, Krasnojarsk, Krasnojarsk, Russian Federation.

4. Ubsunur Biosphere Reserve, Kyzyl, Tyva, Russian Federation.

5. SSAI, Hampton, VA, United States.

6. Climate Science, NASA LaRC, Hampton, VA, United States.

Atmosphere Ocean General Circulations Models (AOGCM) are in agreement that Siberia is expected to experience warming in excess of 40% above global mean temperatures by 2100. Moreover, it is predicted temperature increases will be evident in both the summer and winter. In association with changes in climate, the growing season length, the duration of the fire season, the amount of area burned and fire severity are predicted to increase. Increases in extreme fire seasons are already evident across the circumboreal. Fire will be a major catalyst for ecosystem change, which will force ecosystems to move more rapidly towards a new equilibrium with the climate. Bioclimatic model results predict expansive changes in ecosystems, from a landscape dominated by taiga to a landscape dominated by steppe and forest-steppe.

The focus of this investigation is on south, central Siberia, specifically Tyva, where one would expect to find the initial signs of climate-induced change. The Sayan mountain range offers relatively abrupt change in ecosystems that are often defined by altitude, temperature and precipitation. Tyva is located at a vulnerable southern border, south of the Sayan, and contains 9 Biospheric Reserves, each representing distinct ecosystems. Tyva is home to several relic *Pinus sylvestris* forests.

January temperature increases have exceeded those predicted by the 2090 Hadley Centre scenario, but July temperatures are below predictions. The growing season length has already increased by ~6 to 12 days. Predicted increases in rainfall are not apparent, and generally, precipitation change has been negative. Consequently, several of the relic pine forests have burned (some repeatedly), and natural regeneration is not visible at several sites, even one that had been re-planted on several occasions. Human influences are evident at the local scale (household use) and in some management zones, where forests are logged and planted regardless of their state of regeneration. We present a concentrated and complicated view of one region that is expected to change and may be currently showing indications of climate-induced change.

Contact Information

Amber J. Soja, Hampton, Virginia, USA, 23681-2199, [click here](#) to send an email

ScholarOne Abstracts® (patent #7,257,767 and #7,263,655). © [ScholarOne](#), Inc., 2009. All Rights Reserved.
ScholarOne Abstracts and ScholarOne are registered trademarks of ScholarOne, Inc.
[Terms and Conditions of Use](#)