



Agro-climatic potential across central Siberia in an altered twenty-first century

AGU Poster
GC41C-0832

Tchebakova N.M,* Parfenova E.I.*, Lysanova G.I.** and A.J. Soja***

*Institute of Forest, Siberian Branch, Russian Academy of Sciences, Email: ncbeby@ksc.krasn.ru

**Institute of Geography, Siberian Branch, Russian Academy of Sciences, Email: lysanova@irigs.irk.ru

***National Institute of Aerospace (NIA), Email: Amber.J.Soja@nasa.gov




Fig. 1. The study region Krasnoyarsky Krai and adjacent Republic of Khakassia (hereafter Khakassia) and Republic of Tyva, (hereafter Tyva, bottom). The arable land (brown) currently range across Russia.

Introduction. World concerns about food security are growing as the result of observed and projected climate change impacts on agriculture. The scientific community agrees that climate change consequences differ for agriculture depending on geography; northern countries may benefit from global warming (Russia, northern China), and in the absence of adaptation measures, southern countries and continents (Africa, Australia) may find their food security at risk due to climate warming.

Goals are: to predict 'hot spots' of potential agricultural change in the pre-2010 climate and in the future using Hadley 2020 and 2080 climate change scenarios; and to model future crop potential that may evolve as the climate changes in south-central Siberia, Krasnoyarsk Krai and adjacent Republics of Khakassia and Tyva (Fig.1).




Fig. 2. Observed trends of grain crops, centner ha⁻¹, for 1966-2009 in the forest-steppe zone (green, three farming regions) and steppe zone (red, three farming regions). With warming, crop production increases with sufficient moisture in forest-steppe and reduces as the climate becomes dryer in steppe.

Methods. Our hypothesis is based on previous work (Tchebakova and Parfenova et al.; Soja et al.), which showed the Siberian taiga would shift northwards by the end of the century and at least half of central Siberia would be occupied by steppe and forest-steppe and may become climatically suitable for agriculture. Together with soil potential, suitable deep and fertile soils, these future climates would allow for farming.

Data from 21 weather stations from 1961-2010 in three southern vegetation zones (forest-steppe, steppe, and dry steppe) were used to evaluate climate change trends and climatic indices: growing degree days >5°C and annual moisture index (a ratio between GDD>5°C and annual precipitation). All data were derived from reference books on climate and crops.

Results. From available data, we found that crop output increased during the last half a century (1960-2010) in southern Siberia, (Fig.2). This crop increase may be associated with stable positive trends in both summer and winter temperatures (Fig. 3) that prolonged the growing season up to 1 month over 50 years. Crops decreased in regions where the climate became dryer.

Methods. We constructed simple agri-climatic regression model models that predict the geographic range and yields of annual agri-crops from two summer climatic indices: growing degree-days, GDD, base 5°C, and a moisture index, a ratio between GDD>5°C and annual precipitation.




Fig. 4. Potential climatic ranges (green) of traditional and new crop (italic) species in central Siberia in the 2010 and HadCM3 B1 and A2 2080 climates




Fig.3. January and July temperature trends for 1961-2010 were positive and showed that north of 51° N, January temperatures increased 1-2°C and July temperatures increased 0.7-1.5°C over the last 50 years. In Tyva, both January and July temperature increases were two times greater, 2-4°C and 1.4-3.2°C, respectively. Blue points mark the moisture increase and orange points mark moisture decrease (right).

Results. The crop range and yield models were applied to climatic indices in different time frames: pre-1960; 1960-1990; 1990-2010 (using observed data); and in HadCM3 B1 and A2 2020 and 2080 climates (using climate change scenarios) to predict potential crop transformation for 150 years in a changing climate (Fig. 4 and Fig. 5). Climatically potential crops are restricted by soil potential, which is accepted as limited by sufficiently developed and fertile soils in southern taiga, located north of sub-taiga and forest-steppes. Thus, we use the northern border of the southern taiga as limiting potential agriculture.




Fig. 5. Predicted crop yields (centner ha⁻¹) for traditional grain and maize silage for historic and future periods (Hadley Center climate change scenarios) over the study regions.
Grain: 0. beyond the area of suitable soils; 1. <5; 2.5-10; 3. 10-15; 4. 15-20; 5. 20-25; 6. 25-30; 7. 30-35; 8. > 35;
Silage: 0. beyond the area of suitable soils; 1. <100; 2. 100-200; 3. 200-300; 4. 300-400; 5. 400-500; 6. 500-600; 7. > 600

References.

- * Reference Book on Climate of the USSR 1967-1971 (Leningrad: Gidrometeoizdat)
- * Reference Books on Monthly Climate of Krasnoyarsk Krai, Khakassia, and Tyva 1960-2010 (Novosibirsk: Gidrometeoizdat)
- * Reference Books on Statistical Data. Agriculture of Rep. Khakassia, 1966a-2009 (Abakan: Roscomstat)
- * Reference Books on Statistical Data. Agriculture of Krasnoyarsk Krai 1966b-2009 (Krasnoyarsk: Roscomstat)
- * Reference Books on Statistical Data. Agriculture of Rep. Tyva, 1986-2009 (Kyzyl: Roscomstat)
- * Soja A J, Tchebakova N M, French N H F, Flannigan M D, Shugart H H, Stocks B J, Sukhinin A I, Parfenova E I, Chapin F S III and Stackhouse P W Jr 2007 Climate-induced boreal forest change: Predictions versus current observations *Glob. Plan. Change* **56** 274-96.
- * Tchebakova N M and Parfenova E I 2011 Potential land cover change in Siberia predicted by Siberian bioclimatic model, *10, Environ. Res. Lett.* **6** (2011) 045207
- * Tchebakova N M, Parfenova E I and Soja A J 2009 Effects of climate, permafrost and fire on vegetation change in Siberia in a changing climate *Environ. Res. Lett.* **4** 045013
- * Tchebakova N M, Parfenova E I and Soja A J 2011b Climate change and climate-induced hot spots in forest shifts in central Siberia from observed data *Reg. Environ. Change* (doi: 10.1007/s10113-011-0210-4)
- * Tchebakova N M, Rehfeldt G E and Parfenova E I 2010 From vegetation zones to climatypes: effects of climate warming on Siberian ecosystems *Permafrost Ecosystems. Siberian Larch Forests* ed A Osawa, O A Zyryanova, Y Matsuura, T Kajimoto



A ripe wheat field in Khakassia, southern Siberia

Conclusions.

- 50-85% of central Siberia is predicted to be climatically suitable for agriculture at the end of the century although potential croplands would be limited by the availability of suitable soils within the steppe, forest-steppe, sub-taiga and southern taiga zones;
- Climatic factors control crop distribution and production in southern Siberia (R² = 0.2-0.7);
- Crop production may increase by twofold as climate warms during the century;
- Traditional crops (grain, potato, maize for silage) could gradually shift as far as 500 km northwards (about 50-70 km per decade) and new crops (maize for grain, apricot, grape, gourds) could be introduced in the far south depending on winter conditions and these would necessitate irrigation in the drier 2080 climate.
- Agriculture in central Siberia would likely benefit from climate warming



Acknowledgement. We appreciate the support of the Northern Eurasian Earth Science Partnership Initiative (NEESPI), NASA Land Cover Land Use Change and a NASA Interdisciplinary Science project NNH09ZDA001N-IDS.

